



Topological indices of splitting graph of special case of hexagonal system

M.Bhanumathi¹, K.Easu Julia Rani²

Principal, Govt. Arts College, Kadaladi, Ramanathapuram, India¹

Assistant Professor, Department of Mathematics, T.R.P. Engineering College, Irungalur, Trichy, India²

Abstract: A topological index is a numeric number that helps to find the characteristics of a compounds. There are many applications of graph theory. In this paper we compute first and second Zagreb indices, Randić index, sum-connectivity index, harmonic index, inverse sum indeg index, modified first and second Zagreb indices and first and second hyper-Zagreb indices of the Splitting graph of the special case of hexagonal system.

Keywords: Molecular graph, topological indices, hexagonal system.

I. INTRODUCTION

Topological indices help for the quantitative structures property relationship (QSPR) and quantitative structure activity relationship (QSAR).

Let G be a simple graph, with vertex set $V(G)$ and edge set $E(G)$. The degree d_u of a vertex u is the number of edges that are incident to it.

Since 1947 many number of topological indices have been found. One of the oldest and well known topological indices is the first and second Zagreb indices, was first introduced by Gutman et al. in 1972 [3], and it is defined as

$$M_1(G) = \sum_{uv \in E(G)} [d_G(u) + d_G(v)] \quad \text{and} \\ M_2(G) = \sum_{uv \in E(G)} d_G(u) d_G(v)$$

The connectivity index introduced in 1975 by Milan Randić [4], is defined as

$$R(G) = \sum_{uv \in E(G)} \frac{1}{\sqrt{d_u d_v}}$$

Recently, another variant of the Randić connectivity index called the sum-connectivity index was introduced by B. Zhou and N. Trinajstić [5] in 2008. It is defined as,

$$X(G) = \sum_{uv \in E(G)} \frac{1}{d_u + d_v}$$

In 2014 Jianxi Li and Chee Shiu introduced a new variant of the Randić index named the Harmonic index which first appeared in [6] called the harmonic index $H(G)$ is defined as

$$H(G) = \sum_{uv \in E(G)} \frac{2}{d_u + d_v}$$

Discrete Adriatic indices have been defined by Vukičević and Gašperov in 2010. One among such indices is the inverse sum indeg index, [7] and is defined as

$$ISI(G) = \sum_{uv \in E(G)} \frac{d_u d_v}{d_u + d_v}$$

A. Milicevi, S. Nikoli, N. Trinajstić, introduced in 2004 the modified first and second Zagreb indices [8], and are respectively defined as

$${}^m M_1(G) = \sum_{u \in V(G)} \frac{1}{d_G(u)^2} \\ {}^m M_2(G) = \sum_{uv \in E(G)} \frac{1}{d_G(u) d_G(v)}$$

In 2013 [9], Shirdel et al. introduced the first hyper-Zagreb index of a graph G , which is defined as

$$HM_1(G) = \sum_{uv \in E(G)} [d_G(u) + d_G(v)]^2$$

In [10], the second hyper-Zagreb index of a graph G is defined as

$$HM_2(G) = \sum_{uv \in E(G)} [d_G(u) d_G(v)]^2$$

II. HEXAGONAL SYSTEM

A hexagonal system is defined as a 2-connected plane graph in which every interior face is bounded by a regular hexagon of unit length 1. A hexagonal system H is said to be Catacondensed if it does not possess internal vertices, otherwise H is said to be Pericondensed. A hexagonal chain is a Catacondensed hexagonal system which has no hexagon adjacent to more than two hexagons. Some examples of hexagonal chains can be found in Fig. 1.

One can easily see that, H_{n+1} is a hexagonal chain obtained from H_n by attaching $n+1$ hexagons.

Based on this we define three types of fusion for attaching a new hexagon h_{n+1} to a hexagonal chain H_n with n hexagons $h_1, h_2, h_3, \dots, h_n$:

- If h_{n+1} is on the line 1, it is called α -type fusing;
- If h_{n+1} is on the left-hand side of 1, it is called β -type fusing;
- If h_{n+1} is on the right-hand side of 1, it is called γ -type fusing

where l is the direct line from the center of h_{n-1} to the center of h_n . Any hexagonal chain $H_n (n \geq 2)$ can be obtained from H_2 by a stepwise fusion of new hexagons, and at each step a θ -type fusion is selected, where $\theta \in \{\alpha, \beta, \gamma\}$. Below are examples of the above said three types of fusing in Hexagonal system.




Fig 2.1. Single hexagonal chains with different types of fusions.

III. MAIN RESULTS

In this chapter we compute topological indices of on splitting graph of special case of hexagonal system. That is we are going to study the topological indices of Linear chain with α, β and γ types of fusing. Fig 2 shows the Linear chain with α -type, β -type and γ -type fusing of length 5 which is denoted as $L_5(\alpha, \alpha)$, $L_5(\alpha, \beta)$ and $L_5(\alpha, \gamma)$.




Fig 3.1. Linear chain L_5 with α, β, γ type fusing

Defenition 3.1 [14].

The Splitting graph $S(G)$ of graph G is the graph where for each point v of a graph G , take a new point v' and join v' to all points of G adjacent to v .

The following diagram shows the linear chain $L_3(\alpha, \alpha)$ and its Splitting graph $S[L_3(\alpha, \alpha)]$




Fig 3.2. Linear chain L_3 with α -type fusing and its Splitting graph

Theorem 3.1:

If $S(L_n(\alpha, \alpha))$ is a linear chain with α -type fusing, then

- (i) $M_1(G) = 5(26n + 13)$
 $M_2(G) = 24(11n - 3)$
- (ii) $R(G) = \frac{1}{6}[3(3 + 6\sqrt{2}) + (n - 1)(1 + 2\sqrt{6} + 8\sqrt{3} + 2\sqrt{2})]$
- (iii) $X(G) = \frac{1}{1260}(3465 + 2239(n - 1))$
- (iv) $H(G) = \frac{1}{630}(3465 + 2239(n - 1))$
- (v) $ISI(G) = \frac{1}{35}(980 + 1031(n - 1))$
- (vi) ${}^m M_1(G) = \frac{1}{144}[90(n + 2) + 40(n - 1)]$
 ${}^m M_2(G) = \frac{1}{288}[540 + 280(n - 1)]$
- (vii) $HM_1(G) = 816 + 1158(n - 1)$
 $HM_2(G) = 2304 + 5400(n - 1)$

Proof

Let the graph $G = S(L_n(\alpha, \alpha))$ is a linear chain with α -type fusing as shown in the following diagram




Fig 3.3. Splitting graph of Linear chain L_n with α -type fusing

In $S(L_n(\alpha, \alpha))$ we see that



$$|V(S(L_n(\alpha, \alpha)))| = 4(2n + 1)$$

and $|E(S(L_n(\alpha, \alpha)))| = 3(5n + 1)$. There are 4 types of vertices with degree namely V_2, V_3, V_4 , and V_6 . Hence we have

$$|V_2(S(L_n(\alpha, \alpha)))| = |V_4(S(L_n(\alpha, \alpha)))| = 2(n + 2)$$

and

$$|V_3(S(L_n(\alpha, \alpha)))| = |V_6(S(L_n(\alpha, \alpha)))| = 2(n - 1).$$

Also the edges with degrees of end vertices in $S(L_n(\alpha, \alpha))$ is as follows:

There are $(n - 1)$ edges with degrees of end vertices $(6, 6)$; 6 edges with degrees of end vertices $(4, 4)$; $4(n - 1)$ edges with degrees of end vertices $(4, 6)$; 12 edges with degrees of end vertices $(2, 4)$; $4(n - 1)$ edges with degrees of end vertices $(2, 6)$; $4(n - 1)$ edges with degrees of end vertices $(3, 4)$; $2(n - 1)$ edges with degrees of end vertices $(3, 6)$. Hence

$$(i) \quad M_1(G) = \sum_{uv \in E(G)} [d_G(u) + d_G(v)]$$

$$= (n - 1)(6 + 6) + 6(4 + 4) + 4(n - 1)(4 + 6) + 12(2 + 4) + 4(n - 1)(2 + 6) + 4(n - 1)(3 + 4) + 2(n - 1)(3 + 6) = 5(26n + 13)$$

$$M_2(G) = \sum_{uv \in E(G)} d_G(u) d_G(v) \\ = (n - 1)(6 \times 6) + 6(4 \times 4) + 4(n - 1)(4 \times 6) + 12(2 \times 4) + 4(n - 1)(2 \times 6) + 4(n - 1)(3 \times 4) + 2(n - 1)(3 \times 6) = 24(11n - 3)$$

$$(ii) R(G) = \sum_{uv \in E(G)} \frac{1}{\sqrt{d_u d_v}} \\ = \frac{(n - 1)}{\sqrt{6 \times 6}} + \frac{6}{\sqrt{4 \times 4}} + \frac{4(n - 1)}{\sqrt{4 \times 6}} + \frac{12}{\sqrt{2 \times 4}}$$

$$+ \frac{4(n - 1)}{\sqrt{2 \times 6}} + \frac{4(n - 1)}{\sqrt{3 \times 4}} + \frac{2(n - 1)}{\sqrt{3 \times 6}} = \\ = \frac{1}{6} [3(3 + 6\sqrt{2}) + (n - 1)(1 + 2\sqrt{6} + 8\sqrt{3} + 2\sqrt{2})]$$

$$(iii) X(G) = \sum_{uv \in E(G)} \frac{1}{d_u + d_v} \\ = \frac{(n - 1)}{6 + 6} + \frac{6}{4 + 4} + \frac{4(n - 1)}{4 + 6} + \frac{12}{2 + 4} + \frac{4(n - 1)}{2 + 6} + \frac{4(n - 1)}{3 + 4} + \frac{2(n - 1)}{3 + 6} \\ = \frac{1}{1260} (3465 + 2239(n - 1))$$

$$(iv) H(G) = \sum_{uv \in E(G)} \frac{2}{d_u + d_v}$$

$$\frac{2(n - 1)}{6 + 6} + \frac{2(6)}{4 + 4} + \frac{2(4(n - 1))}{4 + 6} + \frac{2(12)}{2 + 4} \\ + \frac{2(4(n - 1))}{2 + 6} + \frac{2(4(n - 1))}{3 + 4} + \frac{2(2(n - 1))}{3 + 6} \\ = \frac{1}{630} (3465 + 2239(n - 1))$$

$$(v) ISI(G) = \sum_{uv \in E(G)} \frac{d_u d_v}{d_u + d_v} \\ = (n - 1) \left(\frac{6 \times 6}{6 + 6} \right) + 6 \left(\frac{4 \times 4}{4 + 4} \right) + 4(n - 1) \left(\frac{4 \times 6}{4 + 6} \right) + 12 \left(\frac{2 \times 4}{2 + 4} \right) \\ + 4(n - 1) \left(\frac{2 \times 6}{2 + 6} \right) + 2(n - 1) \left(\frac{3 \times 4}{3 + 4} \right) + 2(n - 1) \left(\frac{3 \times 6}{3 + 6} \right) = \frac{1}{35} (980 + 1031(n - 1))$$

$$(vi) {}^m M_1(G) = \sum_{uv \in E(G)} \frac{1}{d_G(u)^2} \\ = 2(n + 2) \left(\frac{1}{2^2} \right) + 2(n + 2) \left(\frac{1}{4^2} \right) + 2(n - 1) \left(\frac{1}{3^2} \right) + 2(n - 1) \left(\frac{1}{6^2} \right) \\ = \frac{1}{144} [90(n + 2) + 10(n + 2) + 4(n - 1) + 1(n - 1)]$$

$${}^m M_2(G) = \sum_{uv \in E(G)} \frac{1}{d_G(u) d_G(v)} \\ = (n - 1) \left(\frac{1}{6 \times 6} \right) + 6 \left(\frac{1}{4 \times 4} \right) + 4(n - 1) \left(\frac{1}{4 \times 6} \right) + 12 \left(\frac{1}{2 \times 4} \right) + 4(n - 1) \left(\frac{1}{2 \times 6} \right) \\ + 4(n - 1) \left(\frac{1}{4 \times 3} \right) + 2(n - 1) \left(\frac{1}{3 \times 6} \right) = \\ = \frac{1}{288} [540 + 280(n - 1)]$$

$$(vii) HM_1(G) = \sum_{uv \in E(G)} [d_G(u) + d_G(v)]^2 \\ = (n - 1)(6 + 6)^2 + 6(4 + 4)^2 + 4(n - 1)(4 + 6)^2 + 12(2 + 4)^2 \\ + 4(n - 1)(2 + 6)^2 + 2(n - 1)(3 + 4)^2 + 2(n - 1)(3 + 6)^2 = 816 + 1158(n - 1)$$

$$HM_2(G) = \sum_{uv \in E(G)} [d_G(u) d_G(v)]^2 \\ = (n - 1)(6 \times 6)^2 + 6(4 \times 4)^2 + 4(n - 1)(4 \times 6)^2 + 12(2 \times 4)^2 \\ + 4(n - 1)(2 \times 6)^2 + 2(n - 1)(3 \times 4)^2 + 2(n - 1)(3 \times 6)^2 = 2304 + 5400(n - 1)$$

Example 3.1:



Consider the graph $G = S(L_4(\alpha, \alpha))$. Then we have, $|V(S(L_4(\alpha, \alpha)))| = 36$; $|E(S(L_4(\alpha, \alpha)))| = 63$. Here $|V_2| = |V_4| = 12$ and $|V_3| = |V_6| = 6$. Also there are 3 edges with degrees of end vertices (6,6); 6 edges with degrees of end vertices (4,4); 12 edges with degrees of end vertices (4,6); 12 edges with degrees of end vertices (2,4); 12 edges with degrees of end vertices (2,6); 12 edges with degrees of end vertices (3,4); 6 edges with degrees of end vertices (3,6). Hence

$$(i) M_1(G) = 510 \\ M_2(G) = 984$$

$$(ii) R(G) = 18.535$$

$$(iii) X(G) = 7.93$$

$$(iv) H(G) = 16.162$$

$$(v) ISI(G) = 116.37$$

$$(vi) {}^m M_1(G) = 4.583$$

$${}^m M_2(G) = 4.792$$

$$(vii) HM_1(G) = 4290$$

$$HM_2(G) = 18504$$

Theorem 3.2:

If $S(L_n(\alpha, \beta))$ is a linear chain with β -type fusing, and $S(L_n(\alpha, \gamma))$ with γ -type fusing then we see that both the chains are symmetric and hence $S(L_n(\alpha, \beta)) = S(L_n(\alpha, \gamma))$. Hence

$$(i) M_1(G) = 130n - 10 \\ M_2(G) = 8(33n - 8)$$

$$(ii) R(G) = \frac{1}{24\sqrt{6}} [6(7\sqrt{6} + 28\sqrt{3} - 12 - 24\sqrt{2}) + 4n(12 + \sqrt{6} + 24\sqrt{2})]$$

$$(iii) X(G) = \frac{1}{2780} (3178980 + 6717n)$$

$$(iv) H(G) = \frac{1}{1890} (1589490 + 6717n)$$

$$(v) ISI(G) = \frac{1}{105} (3093n - 107)$$

$$(vi) {}^m M_1(G) = \frac{1}{144} [90(n+2) + 40(n-1)]$$

$${}^m M_2(G) = \frac{1}{576} [540 + 560n]$$

$$(vii) HM_1(G) = 1158n - 326$$

$$HM_2(G) = 5400n - 2496$$

Proof

Let the graph $G = S(L_n(\alpha, \beta))$ is a linear chain with β -type fusing and $S(L_n(\alpha, \gamma))$ with γ -type fusing as shown in the following diagram




Fig 3.4. Splitting graph of Linear chain L_n with β -type fusing and γ -type fusing

In $S(L_n(\alpha, \beta))$ and $S(L_n(\alpha, \gamma))$ we see that $|V| = 4(2n+1)$ and $|E| = 3(5n+1)$. There are 4 types of vertices with degree namely V_2, V_3, V_4 , and V_6 . Hence we have

$$|V_2| = |V_4| = 2(n+2)$$

$$\text{And } |V_3| = |V_6| = 2(n-1).$$

Also the edges with degrees of end vertices in $S(L_n(\alpha, \beta)) = S(L_n(\alpha, \gamma))$ is as follows:



There are n edges with degrees of end vertices $(6,6)$; 7 edges with degrees of end vertices $(4,4)$; $(4n-6)$ edges with degrees of end vertices $(4,6)$; 14 edges with degrees of end vertices $(2,4)$; $(4n-6)$ edges with degrees of end vertices $(2,6)$; $(4n-6)$ edges with degrees of end vertices $(3,4)$; $2n$ edges with degrees of end vertices $(3,6)$.
Hence

$$(i) M_1(G) = \sum_{uv \in E(G)} [d_G(u) + d_G(v)]$$

$$= n(6+6) + 7(4+4) + (4n-6)(4+6) + 14(2+4) + (4n-6)(2+6) + (4n-6)(3+4) + 2n(3+6)$$

$$= 130n - 10$$

$$M_2(G) = \sum_{uv \in E(G)} d_G(u) d_G(v)$$

$$= n(6 \times 6) + 7(4 \times 4) + (4n-6)(4 \times 6) + 14(2 \times 4) + (4n-6)(2 \times 6) + (4n-6)(3 \times 4) + 2n(3 \times 6) = 8(33n-8)$$

$$(ii) R(G) = \sum_{uv \in E(G)} \frac{1}{\sqrt{d_u d_v}}$$

$$= \frac{n}{\sqrt{6 \times 6}} + \frac{7}{\sqrt{4 \times 4}} + \frac{(4n-6)}{\sqrt{4 \times 6}} + \frac{14}{\sqrt{2 \times 4}} + \frac{(4n-6)}{\sqrt{2 \times 6}} + \frac{(4n-6)}{\sqrt{3 \times 4}} + \frac{2n}{\sqrt{3 \times 6}} = \frac{1}{576} [540 + 560n]$$

$$= \frac{1}{24\sqrt{6}} [6(7\sqrt{6} + 28\sqrt{3} - 12 - 24\sqrt{2}) + 4n(12 + \sqrt{6} + 24\sqrt{2})]$$

$$(iii) X(G) = \sum_{uv \in E(G)} \frac{1}{d_u + d_v}$$

$$= \frac{n}{6+6} + \frac{7}{4+4} + \frac{(4n-6)}{4+6} + \frac{14}{2+4} + \frac{(4n-6)}{2+6} + \frac{(4n-6)}{3+4} + \frac{2n}{3+6}$$

$$= \frac{1}{3780} (3178980 + 6717n)$$

$$(iv) H(G) = \sum_{uv \in E(G)} \frac{2}{d_u + d_v}$$

$$\frac{2n}{6+6} + \frac{2(7)}{4+4} + \frac{2((4n-6))}{4+6} + \frac{2(14)}{2+4} + \frac{2((4n-6))}{2+6} + \frac{2((4n-6))}{3+4} + \frac{2(2n)}{3+6}$$

$$= \frac{1}{1890} (1589490 + 6717n)$$

$$(v) ISI(G) = \sum_{uv \in E(G)} \frac{d_u d_v}{d_u + d_v}$$

$$= n \left(\frac{6 \times 6}{6+6} \right) + 7 \left(\frac{4 \times 4}{4+4} \right) + (4n-6) \left(\frac{4 \times 6}{4+6} \right) + 14 \left(\frac{2 \times 4}{2+4} \right) + (4n-6) \left(\frac{3 \times 4}{3+4} \right) + 2n \left(\frac{3 \times 6}{3+6} \right) = \frac{1}{105} (3093n - 107)$$

$$(vi) {}^m M_1(G) = \sum_{u \in V(G)} \frac{1}{d_G(u)^2}$$

$$= \frac{2(n+2)}{(4n-6)} \left(\frac{1}{24^2} \right) + \frac{2(n+2)}{(4n-6)} \left(\frac{1}{32^2} \right) + \frac{2(n-1)}{(4n-6)} \left(\frac{1}{34^2} \right) + \frac{2(n-1)}{(6^2)} = \frac{1}{144} [90(n+2) + 40(n-1)]$$

$${}^m M_2(G) = \sum_{uv \in E(G)} \frac{1}{d_G(u) d_G(v)}$$

$$= n \left(\frac{1}{6 \times 6} \right) + 7 \left(\frac{1}{4 \times 4} \right) + (4n-6) \left(\frac{1}{4 \times 6} \right) + 14 \left(\frac{1}{2 \times 4} \right) + (4n-6) \left(\frac{1}{2 \times 6} \right) + (4n-6) \left(\frac{1}{3 \times 4} \right) + 2n \left(\frac{1}{3 \times 6} \right) = \frac{1}{576} [540 + 560n]$$

$$(vii) HM_1(G) = \sum_{uv \in E(G)} [d_G(u) + d_G(v)]^2$$

$$= n(6+6)^2 + 7(4+4)^2 + (4n-6)(4+6)^2 + 14(2+4)^2 + (4n-6)(2+6)^2 + (4n-6)(3+4)^2 + 2n(3+6)^2 = 1158n - 326$$

$$HM_2(G) = \sum_{uv \in E(G)} [d_G(u) d_G(v)]^2$$

$$= n(6 \times 6)^2 + 7(4 \times 4)^2 + (4n-6)(4 \times 6)^2 + 14(2 \times 4)^2 + (4n-6)(2 \times 6)^2 + (4n-6)(3 \times 4)^2 + 2n(3 \times 6)^2 = 5400n - 2496$$

Example 3.2: Consider the graph $G = S(L_5(\alpha, \beta))$. Then we have, $|V(S(L_5(\alpha, \beta)))| = 44$; $|E(S(L_5(\alpha, \beta)))| = 78$. Here $|V_2| = |V_4| = 14$ and $|V_3| = |V_6| = 8$. Also there are 5 edges with degrees of end vertices $(6,6)$; 7 edges with degrees of end vertices $(4,4)$; 14 edges with degrees of end vertices $(4,6)$; 14 edges with degrees of end vertices $(2,4)$; 14 edges with degrees of end vertices $(2,6)$; 14 edges with degrees of end vertices $(3,4)$; 2n edges with degrees of end vertices $(3,6)$.



vertices(3,4);10 edges with degrees of end vertices(3,6).
Hence

$$(i)M_1(G) = 640$$

$$M_2(G) = 1136$$

$$(ii)R(G) = 2.830$$

$$(iii)X(G) = 8.775$$

$$(iv)H(G) = 19.772$$

$$(v)ISI(G) = 146.27$$

$$(vi) {}^mM_1(G) = 5.486$$

$${}^mM_2(G) = 5.7986$$

$$(vii)HM_1(G) = 5464$$

$$HM_2(G) = 22712$$

[11] Estrada, E., Torres, L., Rodriguez, L., Gutman, I.: An atom-bond connectivity index: modelling the enthalpy of formation of alkanes. Indian J Chem Sect. A **37**, 849–855 (1998)

[12] Vukicevic, D. and Furtula, B. (2009) Topological Index Based on the Ratios of Geometrical and Arithmetical Means of End-Vertex Degrees of Edges. Journal of Mathematical Chemistry, **46**, 1369-1376

[13]. Furtula, B. Graovac, A. Vukićević, D, Augmented Zagreb index. J. Math. Chem. 48, 370–380 (2010)

[14]. K. Thilakam, A. Sumathi, Wiener Index of Degree Splitting Graph of some Hydrocarbons International Journal of Computer Applications Volume 93 – No 3, May 2014(27-31)

REFERENCES

- [1] Jie Zhang, Hanyuan Deng, The hexagonal chains with the extremal third-order Randić index.
- [2] I. Gutman, A formula for the Wiener number of trees and its extension to graphs containing cycles, Graph Theory Notes New York **27** (1994) 9-15
- [3] I. Gutman, N. Trinajstić, Graph theory and molecular orbitals. Total π -electron energy of alternant hydrocarbons, Chem. Phys. Lett., **17**(1972), 535–538.
- [4] M. Randić, On characterization of molecular branching, J. Amer. Chem. Soc., **97**, (1975), 6609
- [5] D. Vukićević and B. Furtula, Topological index based on the ratios of geometrical and arithmetical means of end-vertex degrees of edges, J. Math. Chem., **46**, (2009), 1369
- [6] S. Fajtlowicz, On conjectures of GraffitiIII, Congr. Number., **60** (1987), pp.187–197.
- [7] D. Vukićević, M. Gaspárov, Bond Additive Modelling 1. Adriatic Indices, Croat. Chem. Acta **83** (2010), 243-260
- [8] S. Nikolić, G. Kovačević, A. Milićević and N. Trinajstić, The Zagreb indices 30 years after, Croatica Chemica Acta **76**(2), (2003) 113-124.
- [9] G.H.Shirdel, H. Rezapour and A.M. Sayadi, The hyper-Zagreb index of graph operations, Iranian J. Math. Chem. **4**(2) (2013) 213-220.
- [10] M.R. Farahani, M.R.R.Kanna and R. Pradeep Kumar, On the hyper-Zagreb indices of nanostructures, Asian Academic Research J. Multidisciplinary, **3**(1), (2016).